6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

Data Sheet
PCSK9-LDLR TR-FRET Assay Kit
Catalog # 72010

DESCRIPTION:
The *PCSK9-LDLR TR-FRET Assay* Kit is designed to measure the inhibition of PCSK9 binding to LDLR in a homogeneous 384 reaction format. PCSK9 is known to function as a negative regulator of hepatic low-density lipoprotein receptors (LDLRs) by binding to the LDLR ectodomain. This FRET-based assay requires no time-consuming washing steps, making it especially suitable for high throughput screening applications. The assay procedure is straightforward and simple; a sample containing europium-labeled (Eu) LDLR ectodomain, dye-labeled acceptor, biotin-labeled PCSK9, and an inhibitor is incubated for two hours. Then, the fluorescence intensity is measured using a fluorescence reader.

COMPONENTS:

Catalog #	Component	Amount	Storage	
79079	LDLR-Eu	2 µg	-80°C	***(Avoid freeze/ thaw cycles!)***
71304	PCSK9, Biotinylated	30 µg	-80°C	
	Dye-Labeled Acceptor	2 x 10 µl	-20°C	
	3x PCSK9 TR-FRET Assay Buffer	4 ml	-20°C	
	White, Nonbinding, low volume, 384-well microtiter plate	1	Room temp.	

MATERIALS REQUIRED BUT NOT SUPPLIED:
Fluorescent microplate reader capable of measuring Time Resolved Fluorescence Resonance Energy Transfer (TR-FRET)
Adjustable micropipettor and sterile tips

APPLICATIONS: Great for screening small molecular inhibitors for drug discovery and HTS applications.

STABILITY: At least 6 months from date of receipt when stored as directed.

REFERENCE(S):
1. Chan, J.C. *et al.* (2009). *Proc. Natl Acad. Sci. USA*, **106**, 9820-9825.
2. Liang, H., *et al.* (2012) *J. Pharmacol. Exp. Ther*. **340** 2289-236.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

ASSAY PROTOCOL:
All samples and controls should be tested in duplicate.

Step 1:

1) Dilute one part **3x PCSK9 TR-FRET Assay Buffer** with 2 parts distilled water (3-fold dilution) to make **1x PCSK9 TR-FRET Assay Buffer**. Make only a sufficient quantity needed for the assay; store remaining stock solution in aliquots at -20°C.

2) Dilute **Dye-Labeled Acceptor** 100-fold in **1x PCSK9 TR-FRET Assay Buffer**. Make only sufficient quantities needed for the assay; store remaining stock solution in aliquots at -20°C.

3) Thaw **LDLR-Eu** on ice. Upon first thaw, briefly spin tube containing **LDLR-Eu** to recover the full contents of the tube. Aliquot into single-use aliquots. Store remaining undiluted **LDLR-Eu** at −80°C immediately. *Note: **LDLR-Eu** is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots.*

4) Dilute **LDLR-Eu** in **1x PCSK9 TR-FRET Assay Buffer** to 0.5 ng/µl. Make only sufficient quantities needed for the assay; store remaining stock solution in aliquots at -20°C.

5) Prepare the master mixture: N wells × (5 µl diluted **Dye-Labeled Acceptor** + 5 µl diluted **LDLR-Eu** + 3 µl **1x PCSK9 TR-FRET Assay Buffer**). Add 13 µl to every well.

6) Add 2 µl of inhibitor solution to each well designated “Test Inhibitor.” Add 2 µl of 10% DMSO in water (inhibitor buffer) to the wells labeled “Negative Control” and “Positive Control.”

7) Add 5 µl **1x PCSK9 TR-FRET Assay Buffer** to wells designated for “Negative Control.”

	Positive Control	**Negative Control**	**Test Inhibitor**
Diluted Dye-Labeled Acceptor	5 µl	5 µl	5 µl
LDLR-Eu (0.5 ng/µll)	5 µl	5 µl	5 µl
1x PCSK9 Assay Buffer	3 µl	8 µl	3 µl
Test Inhibitor	–	–	2 µl
10% DMSO in water (Inhibitor buffer)	2µl	2µl	–
PCSK9, biotinylated (8 ng/µl)	5 µl	0 µl	5 µl
Total	**20 µl**	**20 µl**	**20 µl**

8) Thaw **PCSK9, Biotinylated** protein on ice. Upon first thaw, briefly spin tube containing protein to recover the full contents of the tube. Aliquot **PCSK9, Biotinylated** into single-use aliquots. Store remaining undiluted **PCSK9, Biotinylated** in aliquots at −80°C

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

immediately. *Note:* ***PCSK9, Biotinylated*** *is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots or diluted protein.*

9) Dilute **PCSK9, Biotinylated** in **1x PCSK9 TR-FRET Assay Buffer** to 8 ng/µl. Initiate reaction by adding 5 µl of diluted **PCSK9, Biotinylated** to wells designated for the "Positive Control" and "Test Inhibitor." Discard any remaining diluted PCSK9 protein after use.

10) Incubate at room temperature for 2 hours.

11) Read the fluorescent intensity in a microtiter-plate reader capable of TR-FRET.

Instrument Settings

Reading Mode	Time Resolved
Excitation Wavelength	320±10 nm
Emission Wavelength	620±10 nm
Lag Time	60 µs
Integration Time	500 µs
Excitation Wavelength	320±10 nm
Emission Wavelength	665±10 nm
Lag Time	60 µs
Integration Time	500 µs

CALCULATING RESULTS:

Two sequential measurements should be conducted. Tb-donor emission should be measured at 620 nm followed by dye-acceptor emission at 665 nm. Data analysis is performed using the TR-FRET ratio (665 nm emission/620 nm emission).

If desired, data can be normalized to percent inhibition. Typically for inhibitor screens the FRET value from the positive control is set to zero percent inhibition and the FRET value from the negative control is set to one hundred percent inhibition.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

6042 Cornerstone Court W, Ste B
San Diego, CA 92121
Tel: 1.858.202.1401
Fax: 1.858.481.8694
Email: info@bpsbioscience.com

EXAMPLE OF ASSAY RESULTS:



Figure Legend: Interaction of PCSK9 with LDLR. Data in the above graphs are expressed as FRET ratios. *Data shown is lot-specific. For lot-specific information, please contact BPS Bioscience, Inc. at info@bpsbioscience.com*

RELATED PRODUCTS:

Product	Catalog #	Size
PCSK9, His-tag	71204	50 µg
LDLR, FLAG-tag	71205	50 µg
PCSK9, C-terminal His-Avi, Biotin-labeled	71304	20 µg
LDLR, Biotin-labeled	71206	20 µg
PCSK9 Neutralizing Antibody	71207	50 µg
PCSK9[Biotinylated]-LDLR Binding Assay Kit	72002	96 rxns
PCSK9(D374T)-LDLR TR-FRET Assay Kit	72011	384 rxns

Note: The Dye-Labeled Acceptor used in this assay is a product of Cisbio Bioassays.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.